

Subcounty level estimates and credible intervals for selected reproductive, maternal, newborn, child, and adolescent health and development indicators for 2014 (DHS-7) and 2022 (DHS-8) Kenya and their change over time, version 1.0

25/11/24

Release Content and Descriptions

DHS_7_8_KEN_indicators_and_CI_subcounty.gpkg

DHS_7_8_KEN_indicators_and_CI_subcounty.csv

ANC_4plus

The proportion of women with a live birth in the five years preceding the survey and who had four or more antenatal care visits. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

ANC_blood

The proportion of women with a live birth in the five years preceding the survey who received antenatal care for the most recent birth with blood sample taken. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

ANC_suppl

The proportion of women with a live birth in the five years preceding the survey who received iron tablets or syrup during antenatal care. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution

outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

ANC_timing

The proportion of women who had a live birth in the five years preceding the survey whose first antenatal care visit was at less than 4 months. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

ANC_urine

The proportion of women with a live birth in the five years preceding the survey who received antenatal care for the most recent birth with urine sample taken. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

C_Prev

The proportion of currently married or in union women currently using any modern method of contraception. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

Child_m_15_49

The proportion of women whose first marriage or consensual union occurred before the age of 15 over the full sample of women aged 15-49. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change

between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

Labour_fem

The proportion of currently married or in union women employed in the 12 months preceding the survey. The indicator includes those who worked in the past year, those who were currently working and those who have a job but were on leave over the last 7 days. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

Min_diet

The proportion of children aged 6-23 months who received a minimum acceptable diet. This indicator is a composite of children fed with a minimum dietary diversity and a minimum meal frequency. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

NAR_prim

The proportion of primary school aged children attending primary school. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

NAR_sec

The proportion of secondary school aged children attending secondary school. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

Stunting

The proportion of children under 5 years old stunted (below -2 standard deviations of height-for-age according to WHO standard). The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

Teen_Pregn

Proportion of women 15 to 19 years old who had given birth or were pregnant with their first child over the full sample of women aged 15 to 49. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

U_Pregn

Proportion of births that were either wanted earlier or later than occurred (mistimed) or not wanted at all. The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

Wash_sanit

The proportion of the population with access to improved toilet facilities (Improved sanitation facilities include flush toilet, pit latrine (with at least a slab) or a composting toilet). The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

Wash_water

Percentage (%) of the population with access to improved drinking water (Improved sources of drinking water include water piped into dwelling or yard/plot, to a neighbour, access to a public tap/standpipe, tube well or borehole, protected well, protected spring, rainwater, tanker truck, cart with small tank and bottled water). The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

Wasting

Proportion of children wasted (below to 2 SD of weight-for-height according to the World Health Organisation's (WHO) standard). The subcounty level estimates including mean, lower 95% credible interval (L), upper 95% credible interval (U) are aggregated from high resolution outputs of models using data collected during DHS-7 (R1), DHS-8 (R2), and the difference between both surveys (CH) defined as round 2 - round1. Additionally, for the change between the surveys, an additional variable (P) is included, which represents the probability that the true change in the indicator is greater than 0.

License

These data may be redistributed following the terms of a Creative Commons Attribution 4.0 International (CC by 4.0) license.

Suggested Citation

Dorey, P., Chan, H.M.T, Williams, E.M., Priyatikanto R., Bonnie, A., Tejedor-Garavito, N., Johnson, M., Tatem, A.J. and Pezzulo, C. 2024. Subcounty level estimates and credible intervals for selected reproductive, maternal, newborn, child, and adolescent health and development indicators for 2014 (DHS-7) and 2022 (DHS-8) Kenya, version 1.0 WorldPop, University of Southampton. DOI: <https://doi.org/10.5258/SOTON/WP00791>.

Source Data

This work is based on the Kenya Demographic Health Survey 7 (DHS7) 2014 and the Kenya (DHS8) 2022. The 2014 and 2022 Kenya DHS-7 and DHS-8 were conducted by [1, 2]. Microdata and more information can be found here: <https://dhsprogram.com//> and on relevant Country Reports [1,2].

Indicators were adapted from the open-source code shared by the DHS Program Code Share Project (<https://github.com/DHSProgram>) [3].

Boundary data is taken from Population Division, U.S. Census Bureau. The U.S. Census Bureau's products are open access and can be accessed from <https://www.census.gov/geographies/mapping-files/time-series/demo/international-programs/subnationalpopulation.html>

Population data is from Worldpop Kenya unconstrained population raster 1km resolution for 2014 and 2020. [6]

Methods Overview

We constructed spatial binomial generalised linear models for selected health and development indicators collected from 2014 (DHS-7) and 2022 (DHS-8) Kenya along with geospatial covariates representing geographical, environmental, and socioeconomic factors that are known to influence the indicators. The constructed models are then fitted in the Bayesian framework using the Integrated Nested Laplace Approximation – Stochastic Partial Differential Equations (INLA-SPDE) method [5, 6]. From these models, posterior samples of the grid level estimates (1x1Km) for the whole of Kenya are produced for each indicator at both round 1 (DHS-7) and round 2 (DHS-8) and the change between round 1 and round 2 (round2 – round1). These posterior samples are then aggregated using subcounty boundaries taken from US Census Bureau. The aggregation process weights the grid level posterior samples by population. This aggregation results in posterior samples of estimates for each indicator and time point at subcounty level. From these posterior samples, mean and lower and upper 95%

credible intervals for the estimates at subcounty level are calculated. Additionally for the estimates of the change in the indicator a further variable is produced identifying the probability that the value of the true change is greater than 0. This is calculated as the proportion of the posterior samples which are greater than 0.

The code to produce these outputs is available at <https://doi.org/10.5281/zenodo.14217827>

Acknowledgement

The authors acknowledge the support of the PMO Team at WorldPop. Moreover, the authors would like to thank the DHS Program staff for their input on the construction of some of the indicators. This work was approved by the ethics and research governance committee at the University of Southampton.

References

- 2014 (DHS-7) Kenya DHS: Kenya National Bureau of Statistics, Ministry of Health/Kenya, National AIDS Control Council/Kenya, Kenya Medical Research Institute, National Council for Population and Development/Kenya, and ICF International. 2015. Kenya Demographic and Health Survey 2014 [DATASETS]. Rockville, MD, USA: Kenya National Bureau of Statistics, Ministry of Health/Kenya, National AIDS Control Council/Kenya, Kenya Medical Research Institute, National Council for Population and Development/Kenya, and ICF International.
- 2022 (DHS-8) Kenya DHS: KNBS and ICF. 2023. Kenya Demographic and Health Survey 2022: [DATASETS]. Nairobi, Kenya, and Rockville, Maryland, USA: KNBS and ICF.
- The DHS Program Code Share Project, Code Library, DHS Program, 2022. DHS Program GitHub site. <https://github.com/DHSProgram>.
- Rue, H., Martino, S. and Chopin, N., 2009. Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 71(2), pp.319-392.
- Lindgren, F., Rue, H. and Lindström, J., 2011. An explicit link between Gaussian fields and Gaussian Markov random fields: the stochastic partial differential equation approach. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 73(4), pp.423-498.
- WorldPop (www.worldpop.org - School of Geography and Environmental Science, University of Southampton; Department of Geography and Geosciences, University of Louisville; Département de Géographie, Université de Namur) and Center for International Earth Science Information Network (CIESIN), Columbia University (2018). Global High Resolution Population Denominators Project - Funded by The Bill and Melinda Gates Foundation (OPP1134076). <https://dx.doi.org/10.5258/SOTON/WP00670>