

# Release Statement

Grid level (5x5km) prediction and uncertainty surfaces for selected reproductive, maternal, newborn, child, and adolescent health and development indicators for 2010 (DHS-VI) and 2021 (DHS-8) Burkina Faso and their change over time, version 1.0

29/01/25

## Release Content and Descriptions

<p>Anaemia_round1.tif          Anaemia_round1_sd.tif          Anaemia_round1_lower.tif          Anaemia_round1_median.tif          Anaemia_round1_upper.tif          Anaemia_round2.tif          Anaemia_round2_sd.tif          Anaemia_round2_lower.tif          Anaemia_round2_median.tif          Anaemia_round2_upper.tif          Anaemia_change.tif          Anaemia_change_sd.tif          Anaemia_change_lower.tif          Anaemia_change_median.tif          Anaemia_change_upper.tif</p>	<p>The proportion of women aged 15-49 with mild, moderate or severe anemia or with any anemia. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>ANC_4plus_round1.tif          ANC_4plus_round1_sd.tif          ANC_4plus_round1_lower.tif          ANC_4plus_round1_median.tif          ANC_4plus_round1_upper.tif          ANC_4plus_round2.tif          ANC_4plus_round2_sd.tif          ANC_4plus_round2_lower.tif          ANC_4plus_round2_median.tif          ANC_4plus_round2_upper.tif          ANC_4plus_change.tif          ANC_4plus_change_sd.tif          ANC_4plus_change_lower.tif</p>	<p>The proportion of women with a live birth in the five years preceding the survey and who had four or more antenatal care visits. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>

<p>ANC_4plus_change_median.tif ANC_4plus_change_upper.tif</p>	
<p>ANC_blood_round1.tif ANC_blood_round1_sd.tif ANC_blood_round1_lower.tif ANC_blood_round1_median.tif ANC_blood_round1_upper.tif ANC_blood_round2.tif ANC_blood_round2_sd.tif ANC_blood_round2_lower.tif ANC_blood_round2_median.tif ANC_blood_round2_upper.tif ANC_blood_change.tif ANC_blood_change_sd.tif ANC_blood_change_lower.tif ANC_blood_change_median.tif ANC_blood_change_upper.tif</p>	<p>The proportion of women with a live birth in the five years preceding the survey who received antenatal care for the most recent birth with blood sample taken. The grid level estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change).</p>
<p>ANC_iron_round1.tif ANC_iron_round1_sd.tif ANC_iron_round1_lower.tif ANC_iron_round1_median.tif ANC_iron_round1_upper.tif ANC_iron_round2.tif ANC_iron_round2_sd.tif ANC_iron_round2_lower.tif ANC_iron_round2_median.tif ANC_iron_round2_upper.tif ANC_iron_change.tif ANC_iron_change_sd.tif ANC_iron_change_lower.tif ANC_iron_change_median.tif ANC_iron_change_upper.tif</p>	<p>The proportion of women with a live birth in the five years preceding the survey who received iron tablets or syrup during antenatal care. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>ANC_timing_round1.tif ANC_timing_round1_sd.tif ANC_timing_round1_lower.tif ANC_timing_round1_median.tif ANC_timing_round1_upper.tif</p>	<p>The proportion of women who had a live birth in the five years preceding the survey whose first antenatal care visit was at less than 4 months. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95%</p>

<p>ANC_timing_round2.tif  ANC_timing_round2_sd.tif  ANC_timing_round2_lower.tif  ANC_timing_round2_median.tif  ANC_timing_round2_upper.tif  ANC_timing_change.tif  ANC_timing_change_sd.tif  ANC_timing_change_lower.tif  ANC_timing_change_median.tif  ANC_timing_change_upper.tif</p>	<p>credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>ANC_urine_round1.tif  ANC_urine_round1_sd.tif  ANC_urine_round1_lower.tif  ANC_urine_round1_median.tif  ANC_urine_round1_upper.tif  ANC_urine_round2.tif  ANC_urine_round2_sd.tif  ANC_urine_round2_lower.tif  ANC_urine_round2_median.tif  ANC_urine_round2_upper.tif  ANC_urine_change.tif  ANC_urine_change_sd.tif  ANC_urine_change_lower.tif  ANC_urine_change_median.tif  ANC_urine_change_upper.tif</p>	<p>The proportion of women with a live birth in the five years preceding the survey who received antenatal care for the most recent birth with urine sample taken. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>C_Prev_round1.tif  C_Prev_round1_sd.tif  C_Prev_round1_lower.tif  C_Prev_round1_median.tif  C_Prev_round1_upper.tif  C_Prev_round2.tif  C_Prev_round2_sd.tif  C_Prev_round2_lower.tif  C_Prev_round2_median.tif  C_Prev_round2_upper.tif  C_Prev_change.tif  C_Prev_change_sd.tif  C_Prev_change_lower.tif</p>	<p>The proportion of currently married or in union women currently using any modern method of contraception. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>

<p>C_Prev_change_median.tif C_Prev_change_upper.tif</p>	
<p>Child_m_15_49_round1.tif Child_m_15_49_round1_sd.tif Child_m_15_49_round1_lower.tif Child_m_15_49_round1_median.tif Child_m_15_49_round1_upper.tif Child_m_15_49__round2.tif Child_m_15_49_round2_sd.tif Child_m_15_49_round2_lower.tif Child_m_15_49_round2_median.tif Child_m_15_49_round2_upper.tif Child_m_15_49_change.tif Child_m_15_49_change_sd.tif Child_m_15_49_change_lower.tif Child_m_15_49_change_median.tif Child_m_15_49_change_upper.tif</p>	<p>The proportion of women whose first marriage or consensual union occurred before the age of 15 over the full sample of women aged 15-49. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>Child_m_20_24_round1.tif Child_m_20_24_round1_sd.tif Child_m_20_24_round1_lower.tif Child_m_20_24_round1_median.tif Child_m_20_24_round1_upper.tif Child_m_20_24__round2.tif Child_m_20_24_round2_sd.tif Child_m_20_24_round2_lower.tif Child_m_20_24_round2_median.tif Child_m_20_24_round2_upper.tif Child_m_20_24_change.tif Child_m_20_24_change_sd.tif Child_m_20_24_change_lower.tif Child_m_20_24_change_median.tif Child_m_20_24_change_upper.tif</p>	<p>The proportion of women whose first marriage or consensual union occurred before the age of 18 over women aged 20-24. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>Exclusive_BF_round1.tif Exclusive_BF_round1_sd.tif Exclusive_BF_round1_lower.tif Exclusive_BF_round1_median.tif Exclusive_BF_round1_upper.tif Exclusive_BF_round2.tif Exclusive_BF_round2_sd.tif</p>	<p>The proportion of youngest children under 6 months who are living with their mother who are exclusively breastfed under 6 months of age. The indicator includes those who worked in the past year, those who were currently working and those who have a job but were on leave over the last 7 days. The grid level (5x5km)</p>

<p>Exclusive_BF_round2_lower.tif  Exclusive_BF_round2_median.tif  Exclusive_BF_round2_upper.tif  Exclusive_BF_change.tif  Exclusive_BF_change_sd.tif  Exclusive_BF_change_lower.tif  Exclusive_BF_change_median.tif  Exclusive_BF_change_upper.tif</p>	<p>estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>Labour_fem_round1.tif  Labour_fem_round1_sd.tif  Labour_fem_round1_lower.tif  Labour_fem_round1_median.tif  Labour_fem_round1_upper.tif  Labour_fem_round2.tif  Labour_fem_round2_sd.tif  Labour_fem_round2_lower.tif  Labour_fem_round2_median.tif  Labour_fem_round2_upper.tif  Labour_fem_change.tif  Labour_fem_change_sd.tif  Labour_fem_change_lower.tif  Labour_fem_change_median.tif  Labour_fem_change_upper.tif</p>	<p>The proportion of currently married or in union women employed in the 12 months preceding the survey. The indicator includes those who worked in the past year, those who were currently working and those who have a job but were on leave over the last 7 days. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>Low_BW_round1.tif  Low_BW_round1_sd.tif  Low_BW_round1_lower.tif  Low_BW_round1_median.tif  Low_BW_round1_upper.tif  Low_BW_round2.tif  Low_BW_round2_sd.tif  Low_BW_round2_lower.tif  Low_BW_round2_median.tif  Low_BW_round2_upper.tif  Low_BW_change.tif  Low_BW_change_sd.tif  Low_BW_change_lower.tif  Low_BW_change_median.tif  Low_BW_change_upper.tif</p>	<p>The proportion of live births in the 5 years preceding the survey that weighed at birth reported as less than 2.5 kg. . The indicator includes those who worked in the past year, those who were currently working and those who have a job but were on leave over the last 7 days. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>

<p>Min_diet_round1.tif  Min_diet_round1_sd.tif  Min_diet_round1_lower.tif  Min_diet_round1_median.tif  Min_diet_round1_upper.tif  Min_diet_round2.tif  Min_diet_round2_sd.tif  Min_diet_round2_lower.tif  Min_diet_round2_median.tif  Min_diet_round2_upper.tif  Min_diet_change.tif  Min_diet_change_sd.tif  Min_diet_change_lower.tif  Min_diet_change_median.tif  Min_diet_change_upper.tif</p>	<p>The proportion of children aged 6-23 months who received a minimum acceptable diet. This indicator is a composite of children fed with a minimum dietary diversity and a minimum meal frequency. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>NAR_prim_round1.tif  NAR_prim_round1_sd.tif  NAR_prim_round1_lower.tif  NAR_prim_round1_median.tif  NAR_prim_round1_upper.tif  NAR_prim_round2.tif  NAR_prim_round2_sd.tif  NAR_prim_round2_lower.tif  NAR_prim_round2_median.tif  NAR_prim_round2_upper.tif  NAR_prim_change.tif  NAR_prim_change_sd.tif  NAR_prim_change_lower.tif  NAR_prim_change_median.tif  NAR_prim_change_upper.tif</p>	<p>The proportion of primary school aged children attending primary school. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>NAR_sec_round1.tif  NAR_sec_round1_sd.tif  NAR_sec_round1_lower.tif  NAR_sec_round1_median.tif  NAR_sec_round1_upper.tif  NAR_sec_round2.tif  NAR_sec_round2_sd.tif</p>	<p>The proportion of secondary school aged children attending secondary school. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and</p>

<p>NAR_sec_round2_lower.tif  NAR_sec_round2_median.tif  NAR_sec_round2_upper.tif  NAR_sec_change.tif  NAR_sec_change_sd.tif  NAR_sec_change_lower.tif  NAR_sec_change_median.tif  NAR_sec_change_upper.tif</p>	<p>the difference between both surveys (change) defined as round 2 - round1.</p>
<p>Stunting_round1.tif  Stunting_round1_sd.tif  Stunting_round1_lower.tif  Stunting_round1_median.tif  Stunting_round1_upper.tif  Stunting_round2.tif  Stunting_round2_sd.tif  Stunting_round2_lower.tif  Stunting_round2_median.tif  Stunting_round2_upper.tif  Stunting_change.tif  Stunting_change_sd.tif  Stunting_change_lower.tif  Stunting_change_median.tif  Stunting_change_upper.tif</p>	<p>The proportion of children under 5 years old stunted (below -2 standard deviations of height-for-age according to WHO standard). The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>Teen_Pregn_round1.tif  Teen_Pregn_round1_sd.tif  Teen_Pregn_round1_lower.tif  Teen_Pregn_round1_median.tif  Teen_Pregn_round1_upper.tif  Teen_Pregn_round2.tif  Teen_Pregn_round2_sd.tif  Teen_Pregn_round2_lower.tif  Teen_Pregn_round2_median.tif  Teen_Pregn_round2_upper.tif  Teen_Pregn_change.tif  Teen_Pregn_change_sd.tif  Teen_Pregn_change_lower.tif  Teen_Pregn_change_median.tif  Teen_Pregn_change_upper.tif</p>	<p>The proportion of women between 15-19 years old who have given birth over are pregnant with their first child over the full sample of women aged between 15-49 years old. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>

<p>Wash_sanit_round1.tif  Wash_sanit_round1_sd.tif  Wash_sanit_round1_lower.tif  Wash_sanit_round1_median.tif  Wash_sanit_round1_upper.tif  Wash_sanit_round2.tif  Wash_sanit_round2_sd.tif  Wash_sanit_round2_lower.tif  Wash_sanit_round2_median.tif  Wash_sanit_round2_upper.tif  Wash_sanit_change.tif  Wash_sanit_change_sd.tif  Wash_sanit_change_lower.tif  Wash_sanit_change_median.tif  Wash_sanit_change_upper.tif</p>	<p>The proportion of households with an improved sanitation facility. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>Wash_water_round1.tif  Wash_water_round1_sd.tif  Wash_water_round1_lower.tif  Wash_water_round1_median.tif  Wash_water_round1_upper.tif  Wash_water_round2.tif  Wash_water_round2_sd.tif  Wash_water_round2_lower.tif  Wash_water_round2_median.tif  Wash_water_round2_upper.tif  Wash_water_change.tif  Wash_water_change_sd.tif  Wash_water_change_lower.tif  Wash_water_change_median.tif  Wash_water_change_upper.tif</p>	<p>The proportion of households whose main source of drinking water is an improved source. The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI (round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1.</p>
<p>Wasting_round1.tif  Wasting_round1_sd.tif  Wasting_round1_lower.tif  Wasting_round1_median.tif  Wasting_round1_upper.tif  Wasting_round2.tif  Wasting_round2_sd.tif</p>	<p>The proportion of children under 5 years old who are wasted (below -2 standard deviations of weight-for-height according to WHO standard). The grid level (5x5km) estimates (mean, SD, lower 95% credible interval, median, upper 95% credible interval) were modelled from data collected during DHS-VI</p>



Wasting_round2_lower.tif Wasting_round2_median.tif Wasting_round2_upper.tif Wasting_change.tif Wasting_change_sd.tif Wasting_change_lower.tif Wasting_change_median.tif Wasting_change_upper.tif	(round1), DHS-8 (round 2), and the difference between both surveys (change) defined as round 2 - round1
---	---

## License

These data may be redistributed following the terms of a Creative Commons Attribution 4.0 International (CC by 4.0) license.

## Suggested Citation

Dorey, P., Chan, H.M.T, Williams, E.M., Priyatikanto R., Bonnie, A., Tejedor-Garavito, N., Johnson, M., Tatem, A.J. and Pezzulo, C. 2024. Grid level prediction and uncertainty surfaces for selected reproductive, maternal, newborn, child, and adolescent health and development indicators for 2010 (DHS-VI) and 2021 (DHS-8) Burkina Faso, version 1.0. WorldPop, University of Southampton. DOI: <https://doi.org/10.5258/SOTON/WP00806>

## Source Data

This work is based on the Burkina Faso Demographic Health Survey VI (DHS-VI) 2010 and the Burkina Faso (DHS-8) 2021. The 2010 and 2021 Burkina Faso DHS-VI and DHS-8 were conducted by [1, 2]. Microdata and more information can be found here: <https://dhsprogram.com/> and on relevant Country Reports [1,2]. The datasets [6] were accessed from the DHS spatial repository [7].

Indicators were adapted from the open-source code shared by the DHS Program Code Share Project (<https://github.com/DHSProgram>) [3].

## Methods Overview

We constructed spatial binomial generalised linear models for selected health and development indicators collected from 2010 (DHS-VI) and 2021 (DHS-8) Burkina Faso

along with geospatial covariates representing geographical, environmental and socioeconomic factors that are known to influence the indicators. The constructed models are then fitted in the Bayesian framework using the Integrated Nested Laplace Approximation – Stochastic Partial Differential Equations (INLA-SPDE) method [4, 5]. Grid-level estimates (5x5km resolution surfaces), including the mean, standard deviation, lower 95% credible interval, median, and upper 95% credible interval were calculated using the fitted models.

The code to produce these outputs is available at <https://doi.org/10.5281/zenodo.14217827>

## Acknowledgement

The authors acknowledge the support of the PMO Team at WorldPop. Moreover, the authors would like to thank the DHS Program staff for their input on the construction of some of the indicators. This work was approved by the ethics and research governance committee at the University of Southampton.

## References

1. Institut National de la Statistique et de la Démographie - INSD/Burkina Faso, & ICF International. (2012). Burkina Faso Enquête Démographique et de Santé et à Indicateurs Multiples (EDSBF-MICS IV) 2010. <http://dhsprogram.com/pubs/pdf/FR256/FR256.pdf>
2. Institut National de la Statistique et de la Démographie, & The DHS Program. (2023). Burkina Faso Enquête Démographique et de Santé 2021. <https://www.dhsprogram.com/pubs/pdf/FR378/FR378.pdf>
3. The DHS Program Code Share Project, Code Library, DHS Program, 2021. DHS Program GitHub site. <https://github.com/DHSProgram>.
4. Rue, H., Martino, S. and Chopin, N., 2009. Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 71(2), pp.319-392.
5. Lindgren, F., Rue, H. and Lindström, J., 2011. An explicit link between Gaussian fields and Gaussian Markov random fields: the stochastic partial differential equation approach. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 73(4), pp.423-498.
6. ICF. 2004-2017. Demographic and Health Surveys (various) Burkina Faso Demographic and Health Survey 2010, Burkina Faso Demographic and Health

Survey 2021[Datasets]. Funded by USAID. Rockville, Maryland: ICF [Institut National de la Statistique et de la Démographie]

7. ICF. The DHS Program Spatial Data Repository. Funded by USAID.  
<https://spatialdata.dhsprogram.com/home/>