

Release Statement

Modelled gridded population estimates for Kasai Province in the Democratic Republic of Congo version 4.2.

13 March 2025

Abstract

This data release provides gridded population estimates (at a spatial resolution of 3 arc-seconds, approximately 100-metre grid cells) for Kasai Province in the Democratic Republic of Congo (DRC), including estimates for various age-sex groups. The project team utilized Pre-Distribution Registration Survey (PDRS) data from the National Malaria Control Programme (PNLP), which were collected during anti-malarial campaigns across the DRC. Due to the absence of recent PDRS data for Kasai, we used data from the neighboring province of Kwilu to train our model and made grid-level predictions for Kasai, using geospatial covariates specific to Kasai. The modelling was done using a Bayesian statistical hierarchical modelling framework. The approach facilitated simultaneous accounting for the multiple levels of variability within the data. It also allowed the quantification of uncertainties in parameter estimates. These model-based population estimates can be considered as most accurately representing the year 2023. Although the methods were robust enough to explicitly account for key random biases within the datasets, it is noted that systematic biases, which may arise from sources other than random errors within the observed data collection process, remain.

These data were produced by the WorldPop Research Group at the University of Southampton. This work was part of the GRID3 – Phase 2 Scaling project, with funding from the Gates Foundation (INV-044979). Project partners included GRID3 Inc, the Center for Integrated Earth System Information (CIESIN) within the Columbia Climate School at Columbia University, and WorldPop at the University of Southampton. The final statistical modelling was designed, developed, and implemented by Chris Nnanatu. Data processing was done by Ortis Yankey with additional support from Heather Chamberlain. Project oversight was done by Chris Nnanatu, Attila Lazar, and Andy Tatem. The PDRS data from the malaria insecticide treated net (ITN) distribution campaigns were collected, processed, anonymised, and shared by the PNL and its implementing partners. The settlement extent data was prepared and shared by CIESIN (2024). The data has been clipped to Grid3-CIESIN health area extent (CIESIN, 2025).

The authors followed rigorous procedures designed to ensure that the used data, the applied method and thus the results are appropriate and of reasonable quality. If users encounter apparent errors or misstatements, they should contact WorldPop at release@worldpop.org.

WorldPop, University of Southampton, and their sponsors offer these data on a "where is, as is" basis; do not offer an express or implied warranty of any kind; do not guarantee the quality, applicability, accuracy, reliability or completeness of any data provided; and shall not be liable for incidental, consequential, or special damages arising out of the use of any data that they offer. These data are operational population estimates and are not official government statistics.

RELEASE CONTENT

1. COD_Kasai_province_population_v4.2_gridded.zip
2. COD_Kasai_province_population_v4.2_agesex.zip

LICENSE

These data may be redistributed following the terms of a [Creative Commons Attribution 4.0 International \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/) license.

SUGGESTED CITATION

Nnanatu C., Yankey O., Abbott T. J., Chamberlain H., Lazar A. N., Tatem A. J. 2025. Bottom-up gridded population estimates for Kasai Province in the Democratic Republic of Congo (2023), version 4.2. WorldPop, University of Southampton. doi: <https://dx.doi.org/10.5258/SOTON/WP00789>

FILE DESCRIPTIONS

The projection for all GIS files is the geographic coordinate system WGS84 (World Geodetic System 1984). Kindly note that while this data represents population counts, values contain decimals, i.e. fractions of people. This is because both the input population data and age-sex proportions contain decimals. For this reason, it is advised to aggregate the rasters at a coarser scale. For example, if four grid cells next to each other have values of 0.25 this indicates that there is 1 person somewhere in those four grid cells.

COD_Kasai_province_population_v4_2_gridded.tif

This geotiff raster contains estimates of total population size for each approximately 100-metre grid cell (0.0008333 decimal degrees grid) across Kasai Province. The values are the mean of the posterior probability distribution for the predicted population size in each grid cell. Grid cells with values of 0 represent areas that were mapped as unsettled according to building footprints data.

COD_Kasai_province_population_v4_2_lower.tif

This geotiff raster contains estimates of the lower bound credible interval (2.5% CI) for each grid cell across Kasai Province. The values are the 2.5% posterior probability distribution of the predicted population size in each grid cell. The lower bound estimates cannot be summed across grid cells to produce a lower credible interval measure for a multi-cell area. Grid cells with values of 0 represent areas that were mapped as unsettled according to building footprints data

COD_Kasai_province_population_v4_2_upper.tif

This geotiff raster contains estimates of the upper bound credible interval (97.5% CI) for each grid cell across Kasai Province. The values are the 97.5% posterior probability distribution for the predicted population size in each grid cell. The upper bound estimates cannot be summed across grid cells to produce an upper bound credible interval measure for a multi-cell area. Grid cells with values of 0 represent areas that were mapped as unsettled according to building footprints data.

COD_Kasai_province_population_v4_2_agesex.zip

This zip file contains 40 geotiff rasters at a spatial resolution of 3 arc-seconds (approximately 100-metre grid cells). Each raster provides gridded population estimates for an age-sex group per grid cell across Kasai. We provide 36 rasters for the commonly reported age-sex groupings of sequential age classes for males and females separately. These are labelled with either an “m”(male) or an “f” (female) followed by the number of the first year of the age class represented by the data. “f0” and “m0” are population counts of under 1-year olds for females and males, respectively. “f1” and “m1” are population counts of 1- to 4-year-olds for females and males, respectively. Over 4 years old, the age groups are in five-year bins labelled with a “5”, “10”, etc. Eighty-year-olds and over are represented by the groups “f80” and “m80”. We provide four additional rasters that represent demographic groups often targeted by programmes and interventions. These are “under1” (all females and males under the age of 1), “under5” (all females and males under the age of 5), “under15” (all females and males under the age of 15) and “f15_49” (all females between the ages of 15 and 49, inclusive). These data were produced using age-sex proportions from the 2024 WorldPop Global subnational population pyramids for the DRC. The age-sex proportions are available per a given province. Hence we applied the age-sex proportions for Kasai to the gridded population estimates (COD_Kasai_province_population_v4_2_gridded.tif) to allocate the population to the different age-sex classes.

RELEASE HISTORY

Version 4.2 (13 March 2025)

- This is a major update the data for Kasai Province [doi: 10.5258/SOTON/WP00789] (as described in this release statement).
- This data is released as part of a collection of population estimates for 11 DRC provinces: <https://wopr.worldpop.org/?COD/Population/v4.2>

Version 3.0 (4 January 2022) [doi:10.5258/SOTON/WP00720]

- Original release of the population dataset for the Haut-Katanga, Haut-Lomami, Ituri, Kasai, KasaiOriental, Lomami and Sud-Kivu provinces.

ASSUMPTIONS AND LIMITATIONS

The gridded population data for Kasai were generated using PDRS data collected in 2023 for Kwilu. Due to the lack of recent PDRS data for Kasai, we utilized data from the neighboring Province of Kwilu to train our model and made grid-level predictions for Kasai using geospatial covariates specific to Kasai. Although Kasai and Kwilu are adjacent, their population distributions may differ significantly, potentially introducing variability in our model's accuracy.

These population estimates most likely represent 2023, but because of the different ages of the input data used to build the model, a precise time point cannot be allocated. The PDRS data for Kwilu that was used as the response variable was collected in 2023, while geospatial covariates data were collected from different time periods between 2020 and 2023. Similarly, the CIESIN settlement layers were produced in 2024. The inherent heterogeneity in the temporal alignment of these datasets used in the modelling may introduce uncertainties and potential inaccuracies in the modelling process.

Data on population per household (household size), collected during ITN distribution campaigns, was aggregated to calculate total population count for a given spatial unit. Given that the number of ITNs received by a household is proportional to the household size, there is an incentive for respondents to potentially inflate counts of population per household. The presence of inflated household sizes in the input population data would likely introduce systematic biases in the modelled estimates.

The statistical model produced unrealistically high population estimates for some grid cells relative to the number of building counts. These grid cells are mostly in rural and remote areas and include those with the following coordinates:

(21°4'2.173" E, 6°52'25.928" S), (20°8'49.488" E, 5°35'2.1" S), and (21°24'55.167" E, 6°12'31.064" S). These seemingly overinflated population numbers might be a function of residual errors not accounted for by the model, and it may also be that the PNLP data used as input in these locations could be highly inflated.

The model does not directly account for external factors such as migration, displacement, or sudden demographic changes, which could significantly influence population dynamics. However, the use of recently collected demographic and settlement datasets which capture recent changes in the population distribution/density offers extra layer of advantage. Nevertheless, the estimates may not fully reflect dynamic population shifts occurring beyond the scope of the input data.

Grid cell alignment is based on a mastergrid. Note that this version's (v4.2) mastergrid aligns with version 4.1 but does not align with previous DRC gridded population layers, namely versions v1.0, v2.0, v3.0. We updated the mastergrid in 2024 to ensure grid cell alignment across all new WorldPop data products.

SOURCE DATA

The key datasets used to produce the modelled population estimates are:

PDRS Data

The input population dataset used for the population modelling for Kasai Province was the PDRS malaria bednet campaign data collected for Kwilu Province. The PDRS dataset, which was collected in 2023, provided detailed information on a given household for which a bednet was issued, such as the household size, the number of bednets issued, the number of children in the household, the number of males, and the number of females, among others.

Although the malaria bednet campaign was designed to distribute bednet to every household within the province, a preliminary exploratory data analysis carried out on the PDRS data indicated that some households were not visited during the campaign, while others were not completely covered.

The GPS points of all households within the province were provided in the PDRS data. We implemented population modelling for small spatial units, utilising unofficial boundaries similar to census Enumeration Areas ("pre-EAs"; Qader et al., 2024). The household-level data on population counts was spatially aggregated to these spatial units, by summing the household size data for all GPS points within each pre-EA boundary.

Settlement Data

Settlement data was provided by CIESIN in the form of raster files (CIESIN, 2024). We obtained two different settlement products, namely (i) settlement area, which indicates the area of all buildings whose centroid falls within a given cell, and (ii) building count,

which is the number of building centroids within a given cell. Each of these settlement layers was used in separate analyses together with the observed population count and ancillary geospatial data in robust statistical modeling. After using each settlement layer in the analysis, we compared model metrics and the gridded population layer from both layers. Settlement building count provided more realistic population numbers at the gridcell level and hence was retained for the final population predictions.

Geospatial Covariates

A wide variety of geospatial covariates, which are related to population distribution, were considered in the modelling. These geospatial covariates include land uses and land cover data, climate variables such as temperature and rainfall, physical features and infrastructure such as roads and schools, and conflict data. Population model covariates were selected using a generalized linear model (GLM) – based stepwise selection method. The selected covariates were further accessed for multi-collinearity and statistical significance. Eventually, of the 85 geospatial covariates initially tested, 10 were retained as the best fit covariates with variance inflation factor (VIF) of less than 5. The descriptions of these final geospatial covariates are presented in Table 1 below.

Table 1. Selected geospatial covariates for the modelling.

Description	Source	Link/Reference
Slope	SRTM	https://www.viewfinderpanoramas.org/dem3.html
Euclidean distance to Grid3 health facilities	GRID3	https://data.grid3.org/datasets/8a8d510bd9404212864348010112212b_0/explore
Euclidean distance to cropland/natural vegetation landcover 2020	WorldPop	Woods et al (2024)
Dry Matter Productivity	Copernicus	https://land.copernicus.eu/global/products/ba
Euclidean distance to OSM waterbodies	OSM	https://www.openstreetmap.org
Euclidean distance to hospitals facilities	WorldPop	Woods et al (2024)
Mean – Normalized vegetation Index 2021	Copernicus	https://land.copernicus.eu/global/products/ba
Euclidean distance to OSM local Roads 2023	OSM	https://www.openstreetmap.org
Euclidean distance to tree covers 2020	WorldPop	Woods et al (2024)
Coefficient of variation – Microsoft building length	Microsoft	https://github.com/microsoft/RoadDetections

Age-Sex Proportions

We used the 2024 WorldPop Global subnational population pyramids (Bondarenko et al 2025) to calculate the age-sex proportions for Kasai. We multiplied our gridded population estimates (COD_Kasai_province_population_v4_2_gridded.tif) by the age-sex proportions(grouping) to produce COD_Kasai_province_population_v4.2_agesex.zip.

METHODS OVERVIEW

The key steps of our approach were as follows:

- Cleaning and summarizing the household sizes from the PDRS dataset to get the total population at the pre- enumeration area (pre-EA) level (Qader et al. 2024). PDRS data points with household sizes above 500 people per household signalled potential outliers and as such we imputed these household sizes with the median household size. Similarly, PDRS data point with household sizes of 0 were also imputed using the median household size
- Geospatial covariates were subjected to robust covariate selection for model training and parameter estimation.
- We developed a hierarchical Bayesian statistical model using the INLA-SPDE approach (Lindgren et al. 2011) to fit and predict the population count.
- Population estimates were predicted at grid cell level using the grid cell values of the covariates selected at the model training level.

Statistical Modelling

We approached the population modelling using two complementary methods. The first method assumes that the PDRS data provides unbiased estimates of the population counts and does not require any form of bias correction (here after known as *unscaled model approach*). Whereas the second method acknowledges the fact that the PDRS data, which was our population model's primary input, could be systematically biased and requires bias adjustment to avoid under- or over-estimation of population counts. Specifically, for the second method, we took advantage of an existing recent Microcensus data (Flowminder Foundation et. al., 2021; UCLA and Kinshasa School of Public Health, 2018) that overlapped with the PDRS data at cluster unit levels across six provinces in the DRC (Kongo Central, Kinshasa, Kwilu, Haut-Lomami, Sud-Kivu, and Ituri). Thus, the second approach which is also known as the *scaled model approach* uses a scaling factor to adjust for the potential bias in the population count. Further details of the models' specifications are provided below:

Unscaled Model Approach

In general, within the context of bottom-up population modelling (Leasure et al. 2022, Boo et al., 2022; Darin et al., 2022, Nnanatu et al. 2022), the observed population count at area unit k , y_k , is a Poisson distributed random variable with mean parameter $\lambda_k = \bar{d}_k B_k$ where k is the estimation unit (e.g., enumeration area), while \bar{d}_k and B_k are the mean parameter of the corresponding population density and the number of buildings/settled area, respectively. That is,

$$y_k \sim \text{Poisson}(\bar{d}_k B_k) \quad (1)$$

Then, the transformed mean population density \bar{d}_k is assumed to be linked to a set of geospatial covariates with log-link function:

$$\log(\bar{d}_k) = \mu + \sum_{j=1}^J \beta_j x_{kj} + \sum_{l=1}^L f_l(z_{kl}) \quad (2)$$

where μ is the intercept parameter, $\boldsymbol{\beta} = (\beta_1, \dots, \beta_J)$ is a vector of fixed effects coefficients of the (x_1, \dots, x_J) geospatial covariates; $f_l(\cdot)$ is a function of L random effects covariates including those that capture variability in the population estimates due to settlement type, cluster location and spatial autocorrelations. The population density (defined as people per building or people per settled area) is assumed to be a Gamma distributed random variable with parameters α and γ with mean and variance given by $\bar{d}_k = \alpha/\gamma$ and $\sigma_d^2 = \alpha/\gamma^2$, respectively.

The inclusion of spatial autocorrelation requires the use of computationally efficient statistical modelling software. Thus, the integrated nested Laplace approximation (INLA; Rue et al 2009; Lindgren et al., 2011) is used via the R-INLA statistical package. Note that the method described above predicts population count at regular grid cells using the parameter values trained at the cluster/pre-EA level by calculating the predicted grid-cell level population density as

$$\hat{d}_g = \exp\left(\hat{\mu} + \sum_{j=1}^J \hat{\beta}_j x_{gj} + \sum_{l=1}^L \hat{f}_l(z_{gl})\right) \quad (3)$$

where $\{x_g\}_{g=1}^G$ are the corresponding grid cell level values of the geospatial covariates used in training the model at the cluster level, so that the overall predicted population count across the G 100m by 100m grid cells is given by

$$\widehat{pop} = \sum_{g=1}^G B_g \hat{d}_g \quad (4)$$

where B_g is the corresponding building count or the size of settled area in grid g . We assumed default INLA priors for each of the parameter estimates which have been found to be robust.

Scaled Model Approach

For the second method, a scale factor is calculated for each of the overlapping clusters across the 6 provinces according to equation (5):

$$c_k = \frac{m_k}{p_k} \quad (5)$$

where c_k , m_k and p_k are the scale/adjustment factor, Microcensus population count, and PDRS population count for cluster k , respectively. That is, when $c_k < 1$, the PDRS overestimates the population count by $100(1 - c_k)\%$, and underestimates by $100(c_k - 1)\%$ when $c_k > 1$. Thus, the first approach described above assumes that the PDRS data is as good as the Microcensus data, that is, $c_k = 1$ (or $m_k = p_k$).

We predicted estimates of the scale factor c_k at locations where Microcensus and PDRS datasets did not overlap using model parameters trained with the scale factor data $\mathbf{c} = (c_1, \dots, c_n)^T$ obtained at the n Microcensus-PDRS overlapping clusters. A similar approach was also adopted to predict the scale factors at grid cell levels to better capture local variations. To do this, first we assume that the scale factors are realisations from Gamma distributed random variables, that is,

$$C_k \sim \text{Gamma}(\alpha_c, \gamma_c) \quad (6)$$

with mean and variance given by $\bar{c}_k = \alpha_c / \gamma_c$ and $\sigma_c^2 = \alpha_c / \gamma_c^2$, respectively. So that the mean parameter is linked to a set of geospatial covariates as well as spatial autocorrelations and other random effects through a log-link function defined in equation (7):

$$\log(\bar{c}_k) = \mu + \sum_{j=1}^J \beta_j x_{kj} + \sum_{l=1}^L f_l(z_{kl}) \quad (7)$$

where μ is the intercept parameter, $\boldsymbol{\beta} = (\beta_1, \dots, \beta_J)$ is a vector of fixed effects coefficients of the (x_1, \dots, x_J) geospatial covariates (See Table 1 for the description of the final covariates selected through stepwise regression); $f_l(\cdot)$ is a function of L random effects covariates including those that capture variability in the population estimates due to settlement type, cluster location and spatial autocorrelations. Then, grid cell predictions of the scale factors \hat{c}_g are obtained using the grid cell values of the geospatial covariates and the estimates of the model parameters $\hat{\mu}$, $\hat{\boldsymbol{\beta}}$ and \hat{f} trained at the cluster level:

$$\hat{c}_g = \exp \left(\hat{\mu} + \sum_{j=1}^J \hat{\beta}_j x_{gj} + \sum_{l=1}^L \hat{f}_l(z_{gl}) \right) \quad (8)$$

Finally, the grid cell level scaled population estimates are obtained as

$$\widehat{pop}_g^{(scaled)} = \hat{c}_g \times \widehat{pop}_g \quad (9)$$

where $\widehat{pop}_g = B_g \hat{d}_g$ is the corresponding unscaled grid cell level population estimate.

In this study, we approached the population modelling through five key steps:

- Fit Bayesian Hierarchical regression model to Kwilu and Kasai data combined to predict Kasai input population data based on their shared boundary and settlement types characteristics only (random intercept geostatistical model adjusting for settlement type and spatial autocorrelation).
 - This was implemented within INLA by joining both datasets together and setting PNLP data values of Kasai to NA
- Train geostatistical model to predict scale factor across the entire DRC provinces.
 - The scale factor is defined as the ratio of the PNLP versus Microcensus observations obtained across all overlapping clusters across DRC provinces.
 - It is used as a correction factor for potentially systematically biased PNLP observations.
- Predict scale factor values for Kasai Province using the trained scale factor model parameters.
- Apply scaling to the predicted input PNLP-based population data for Kasai by multiplying it with the corresponding predicted scale factors. This yields a scaled input data for Kasai, that is, bias corrected input data.
- Train Bayesian Hierarchical regression model using population density based on the scaled input data for Kasai input data with a set of best fit geospatial covariates.
- Finally, use the trained model parameters to predict population numbers at the grid cells.

The novelty of the modelling approach utilised here is that it allows for the adjustment of potential systematic bias in the input population data within a coherent Bayesian hierarchical population modelling framework while at the same time adjusting for spatial autocorrelation within the observed data.

All data processing and analysis was carried out using R (v.4.2.2) (R Core Team, 2023) and INLA (v 22.05.07) (Rue et al. 2009). The concept of bottom-up population modelling for estimating population in the absence of recent census data was described by Leasure et al. (2020). Approaches similar to the one used here for Kasai have been carried out for Papua New Guinea (WorldPop and NSO PNG, 2022) and Cameroun (Nnanatu et al, 2022).

ACKNOWLEDGEMENTS

We thank the DRC PNLP and its implementing partners for providing access to the anonymised household data collected during malaria ITN distribution campaigns, in accordance with the relevant data sharing agreements. The whole WorldPop group are acknowledged for overall project support. We thank Attila Lazar and Heather Chamberlain for reviewing the data and providing thoughtful suggestions prior to this release.

WORKS CITED

- Bondarenko M., Priyatikanto R., Tejedor-Garavito N., Zhang W., McKeen T., Cunningham A., Woods T., Hilton J., Cihan D., Nosatiuk B., Brinkhoff T., Tatem A., Sorichetta A. (2025) Constrained estimates of 2015-2030 total number of people per grid square broken down by gender and age groupings at a resolution of 3 arc (approximately 100m at the equator) R2024B version v1. Global Demographic Data Project - Funded by The Bill and Melinda Gates Foundation (INV-045237). WorldPop - School of Geography and Environmental Science, University of Southampton. DOI:10.5258/SOTON/WP00805
- Boo, G., Darin, E., Leasure, D. R., Dooley, C. A., Chamberlain, H. R., Lázár, A. N., ... & Tatem, A. J. (2022). High-resolution population estimation using household survey data and building footprints. *Nature communications*, 13(1), 1330.
- Center for Integrated Earth System Information (CIESIN), Columbia University, Ministère de la Santé Publique, Hygiène et Prévention, Democratic Republic of the Congo, and GRID3. 2025. GRID3 COD - Health Areas v4.0. New York: Columbia University. <https://doi.org/10.7916/nnew-da26>. Accessed 16 March, 2025.
- Center for International Earth Science Information Network (CIESIN), Columbia University. 2024. GRID3 COD - Settlement Extents v3.0 alpha. Unpublished.

- Darin, E., Kuépié, M., Bassinga, H., Boo, G., Tatem, A. J., & Reeve, P. (2022). The Population Seen from Space: When Satellite Images Come to the Rescue of the Census. *Population*, 77(3), 437-464.
- Flowminder Foundation, École de Santé Publique de Kinshasa (ESPK), WorldPop (University of Southampton), Bureau Central du Recensement (BCR). 2021. Microcensus survey in the provinces of Haut-Katanga, Haut-Katanga, Ituri, Kasaï, Kasaï-Oriental, Lomami, and Sud-Kivu (Democratic Republic of the Congo). Version 1.5. [Dataset].
- INS (2017), *Enquête par grappes à indicateurs multiples, 2017-2018, rapport de résultats de l'enquête. Kinshasa, République Démocratique du Congo.*
- Leasure, D.R., Jochem, W.C., Weber, E.M., Seaman, V., & Tatem, A.J. (2020). High resolution population mapping with limited survey data: a hierarchical Bayesian modelling framework to account for uncertainty. *Proceedings of the National Academy of Sciences of the United States of America*, 117(39): 24173–24179.
- Lindgren, F., Rue, H., & Lindström, J. (2011). An explicit link between Gaussian fields and Gaussian Markov random fields: The stochastic partial differential equation approach. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(4), 423–498
- Nnanatu C.C., Yankey O., Abbott T. J., Lazar A. N., Darin E., Tatem A. J. 2022 Bottom-up gridded population estimates for Cameroon (2022), version 1.0. <https://dx.doi.org/10.5258/SOTON/WP00784>
- Qader S H, Batana Y. M., Kosmidou-Bradley W., Skoufias E., Tatem A. J. 2024. Automatic pre-Enumeration Areas (pre-EAs) delineation and national sampling frame for the Democratic Republic of Congo. Policy Research Working Paper; No. (under review). [DRC - Automatic Pre-Enumeration Area Delineation for National Sample Frame Data Report | Data Catalog \(worldbank.org\)](https://www.worldbank.org/publications/prwp/2024/04/24/drc-automatic-pre-enumeration-area-delineation-for-national-sample-frame-data-report)
- R Core Team. 2023. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org>.
- Rue, H., Martino, S., & Chopin, N. 2009. Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *Journal of the royal statistical society:Series b (statistical methodology)*, 71(2), 319-392

UCLA-DRC Health Research and Training Program (University of California, Los Angeles) and Kinshasa School of Public Health. 2017 and 2018. Kinshasa, Kongo Central and former Bandundu microcensus survey data.

D. Woods, T. McKeen, A. Cunningham, R. Priyakanto, A. Soricheta , A.J. Tatem and M. Bondarenko. 2024 "WorldPop high resolution, harmonised annual global geospatial covariates. Version 1.0" University of Southampton: Southampton, UK. DOI:10.5258/SOTON/WP00772

WorldPop and National Statistical Office of Papua New Guinea. 2022. Census-independent population estimates for Papua New Guinea (2020-21), version 1.0. WorldPop, University of Southampton. DOI: 10.5258/SOTON/WP00763